

CLAIMS

We claim :

1. Method of inhibiting in a mammal formation of
5 neutralizing antibodies directed against an heterologous
protein comprising the step of co-administering to said
mammal, an agent in an amount sufficient to deplete or
inhibit at least some antigen presenting cells of said
mammal, and said heterologous protein and/or a nucleic
10 acid sequence encoding said heterologous protein, said
agent being administered prior or simultaneously to said
heterologous protein and/or a nucleic acid sequence,
thereby inhibiting the production of neutralizing
antibodies against said heterologous protein.

15

2. Method according to claim 1, wherein said agent
is selected among viruses, liposomes, antibodies,
parasites, bacteriae, funguses and or fragments thereof,
and nucleic acid sequence encoding said heterologous
20 protein.

3. Method according to claim 2, wherein said virus
is selected among adenovirus, adenovirus associated
virus, retrovirus, pox virus, vaccinia virus, or
25 fragments thereof.

4. Method according to claim 3, wherein said
adenovirus is selected among wild type human adenovirus
and recombinant adenovirus, or a fragment thereof.

30

0920002, 000301

Rous sarcoma virus long terminal repeat promoter (RSV LTR), myeloproliferative sarcoma virus long terminal repeat (MPSV LTR), simian virus 40 early promoter (SV40 IEP), major late promoter of the adenovirus.

5

11. Method according to claims 1 to 10, further comprising the step of administering to said mammal additional agent to enhance the depletion and/or the inhibition of at least some antigen presenting cells of said mammal.

10

12. Method of inhibiting in a mammal formation of neutralizing antibodies directed against an heterologous protein comprising the step of administering to said mammal a recombinant adenovirus, the genome of which comprising at least a nucleic acid sequence encoding said heterologous protein and regulation sequences, in an amount sufficient to deplete or inhibit at least some antigen presenting cells of said mammal, thereby inhibiting the production of neutralizing antibodies against said heterologous protein.

15

20

13. Method according to claim 12, further comprising the step of administering to said mammal additional adenovirus or a fragment thereof, the genome of which not expressing said heterologous protein, thereby enhancing the amount of adenoviruses to deplete or inhibit at least some antigen presenting cells of said mammal.

25

30

0502002-000301

14. Method according to claims 12 to 13, wherein said mammal is a mouse and wherein the amount of adenovirus particules administered to deplete or inhibits at least some antigen presenting cells of said mouse is equal or greater to 4.10^{10} particles, said particles comprising optionally said additional adenovirus.

15. Method according to claim 14, wherein the amount of adenovirus particules administered to deplete or inhibits at least some antigen presenting cells of said mouse is equal or greater to 6.10^{10} particles.

16. Method according to claims 14 and 15, wherein the amount of said recombinant adenovirus able to form plaque, is equal or greater to 4.10^9 pfu/mouse.

17. Method of producing transgenic mammal expressing an heterologous protein said method comprising the step of inhibiting in said mammal formation of neutralizing antibodies directed against said heterologous protein by the use of the method of claims 1 to 16 thereby allowing a long-lasting expression of said heterologous protein.

18. Method according to claim 17, wherein the mammal is selected among mouse, rat, rabbit, cow, pig, goat, sheep.

19. Method for reducing an anti-heterologous protein immune response in a mammal, including human, subject to

the administration of said heterologous protein and/or nucleic acid sequence encoding said heterologous protein, said method comprising the step of inhibiting in said mammal formation of neutralizing antibodies directed against said heterologous protein by the method of claims 1 to 16.

20. Method according to claim 19, wherein said method is a step of a gene therapy protocol for the treatment of human afflicted with a disease selected among inherited or acquired genetic diseases, infectious diseases, inflammatory diseases, autoimmune diseases, cancers, and the associated syndromes thereof.

21. Method for the therapy of a mammal, including humans, afflicted with a disease characterized by the altered expression of an endogenous protein, said method comprising the step of administering to said mammal said protein and/or nucleic acid sequence encoding said protein, and simultaneously or previously, the step of inhibiting in said mammal formation of neutralizing antibodies directed against said protein by the method of claims 1 to 16.

22. Method according to claims 20 and 21, further comprising the step of co-administering simultaneously, separately or sequentially, to said mammal at least one immune modulators selected among cyclosporin, cyclophosphamide, FK506, desoxyspergualine, interleukin-4, interleukin-12, interferon-gamma, anti-CD4 monoclonal

09920902.000301

antibody, anti-CD8 monoclonal antibody, anti-LFA1 monoclonal antibody, antibody directed against CD40 ligand or CTLA4Ig.

5 23. Method of modulating in a mammal formation of neutralizing antibodies directed against an heterologous protein, said method comprising the steps of :

- 10 (i) Optionally, co-administering to a first mammal, at least one agent and said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein, said agent being administered simultaneously, sequentially or separately with said heterologous protein and/or nucleic acid sequence, and determining at least one amount of said heterologous protein and said agent, sufficient to trigger an immune response against said heterologous protein by said first mammal; optionally, re-performing step (i) until said amount is determined;
- 15 (ii) co-administering to a second mammal said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein, in an amount sufficient to trigger an immune response against said heterologous protein, as determined at step (i) and prior or simultaneously, said agent, in an amount greater than the one determined at step (i) and sufficient to trigger an immune response against said agent and sufficient to deplete or inhibit at least some antigen presenting cells of said mammal, and determining for said second mammal
- 20 24. Method of modulating in a mammal formation of neutralizing antibodies directed against an heterologous protein, said method comprising the steps of :
- 25 (i) Optionally, co-administering to a first mammal, at least one agent and said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein, said agent being administered simultaneously, sequentially or separately with said heterologous protein and/or nucleic acid sequence, and determining at least one amount of said heterologous protein and said agent, sufficient to trigger an immune response against said heterologous protein by said first mammal; optionally, re-performing step (i) until said amount is determined;
- 30 (ii) co-administering to a second mammal said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein, in an amount sufficient to trigger an immune response against said heterologous protein, as determined at step (i) and prior or simultaneously, said agent, in an amount greater than the one determined at step (i) and sufficient to trigger an immune response against said agent and sufficient to deplete or inhibit at least some antigen presenting cells of said mammal, and determining for said second mammal

09920902 "000001"

at least one amount of said agent that reduces and/or suppresses the anti-heterologous protein immune response in said mammal; re-performing step (ii) until said amount is determined; and

5 wherein,

- (a) when one administers to a third mammal, said agent in an amount equal or greater than the one determined at step (i) but lesser than the one determined at step (ii), said mammal produces neutralizing antibodies against said heterologous protein and optionnally against said agent; and
- 10 (b) when one administers to said mammal said agent in an amount equal or greater than the one determined at step (ii), said mammal produces neutralizing antibodies against said agent but produces no or few neutralizing antibodies against said heterologous protein.
- 15

24. Method according to claim 23, wherein an additional agent is further administered to said mammal in step (i) and (ii).

20

25. Method according to claims 23 and 24, wherein the amount of said agent of step (ii) is at least twice the amount of said agent determined at step (i).

25

26. Method according to claims 23, 24, 25, wherein said mammal is a mouse and said agent is an adenovirus, and wherein said agent and said nucleic acid sequence encoding said heterologous protein are simulatenously

30

009204002.000301

co-administered as a recombinant adenovirus, the genome of which comprising at least said nucleic acid sequence encoding said heterologous protein and wherein :

- the amount of said recombinant adenovirus particles of step (i) that triggers an immune response towards said heterologous protein in said mouse without depleting or inhibiting at least some antigen presenting cell of said mouse is below 4.10^{10} particles, and/or the amount of said adenovirus particles able to form plaque is below 4.10^9 pfu/mouse; and
- the amount of said recombinant adenovirus particles of step (ii) that reduces or suppresses the anti-heterologous protein immune response in said mouse is at least equal or greater than 4.10^{10} particles and/or the amount of said adenovirus particles able to form plaque is equal or greater than 4.10^9 pfu/mouse.

27. Use of a method according to claim 23 of inhibiting in a mammal formation of neutralizing antibodies directed against an heterologous protein, said method comprising the step of co-administering to said mammal, said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein and prior or simultaneously said agent in an amount equal or greater than the one determined at step (ii).

28. Use of a method according to claim 23 of triggering in a mammal formation of neutralizing

0920002-00001

antibodies directed against an heterologous protein, said method comprising the step of co-administering simultaneously, separately or sequentially to said mammal said heterologous protein and/or a nucleic acid sequence encoding said heterologous protein, and said agent in an amount and/or concentration equal or greater than the one determined at step (i) but lesser than the one determined at step (ii).

29. Method for the therapy of a mammal affected by a disease wherein at least one endogenous protein is involved in said disease ethiology, said method comprising the step of inhibiting the biological functions of said endogenous protein by enhancing the production of neutralizing antibodies against said protein by use the method according to claim 23.

30. Method according to claim 29, wherein said disease is chosen among auto-immune diseases, inflammatory diseases, cancers, viral infections, bacterial infections, parasites infections, funguses infections.

31. Use of a method according to claim 28 to produce a mammal with a functional inactivation of at least one endogenous protein, said method comprising the step of administering to a mammal in a simultaneous, separate or sequential manner at least one agent and an heterologous protein and/or a nucleic acid sequence encoding for said heterologous protein, said nucleic acid sequence being

09920902.000301

administering to said mammal in a simultaneous, separate or sequential manner, at least one agent and said heterologous protein and/or a nucleic acid sequence encoding for said heterologous protein, said nucleic acid sequence being expressed in at least one cell of said mammal, wherein the amount of said heterologous protein, optionally of said agent, is at least sufficient to trigger an immune response against said heterologous protein and the amount of said agent is not sufficient to deplete or inhibit at least some antigen presenting cells of said mammal.

36. Mammal obtained by the method of claim 35.

37. Mammal of claim 36, wherein said mammal is a mouse, rat, rabbit.

38. Use of a mammal according to claims 36 and 37 to perform biological, physiological, biochemical, molecular studies and analysis of the function of said heterologous and/or homologous protein.

39. Use of a mammal according to claims 36 and 37 to perform drug screening.

40. Use of a mammal according to claims 36 and 37 to isolate spleen cells from said mammal that expresses antibody directed against said heterologous an/or endogenous protein to make hybridoma(s).

41. Use of biological fluid of the mammal according to claims 26 and 27 to prepare serum and/or polyclonal antibodies.

10 43. Method according to claims 1 to 26 and 33, use
of a method according to claim 27 to 32, wherein said
heterologous protein or a fragment thereof is selected
among the proteins that are presented by class I major
15 histocompatibility molecule (CMH I), a class II major
histocompatibility molecule (CMH II), or a combination
of a class I major histocompatibility molecule and a
class II major histocompatibility molecule.

25 45. Method according to claim 44, wherein said
secreted protein is selected among neuromediators,
hormones, interleukines, lymphokines, interferons,
chemokines, growth factors.

